

Total Shoulder Arthroplasty Rehabilitation Post-Operative Guidelines

Total shoulder arthroplasty (TSA) is often indicated for end-stage osteoarthritis, rheumatoid arthritis when rotator cuff is viable, avascular necrosis with glenoid chondral wear, and acute fracture or posttraumatic arthritis with intact rotator cuff. Pre-operative pathology will impact functional recovery as range of motion tends to be better following TSA for primary osteoarthritis compared to arthroplasty for rheumatoid arthritis, posttraumatic arthritis and rotator cuff arthropathy. Most patients will achieve 90-100% functional improvement by 6 months though pre-operative status, co-morbidities and specifics of surgical procedure/approach can impact outcome. It is imperative to review operative notes and communicate with surgeon, as repair of either subscapularis or long head of biceps may have occurred, requiring additional protection.

Listed below is a suggested TSA rehabilitation guideline. Please confirm with the surgeon for specific precautions and guidelines if necessary (**will reference physician preferences*).

Patient Information:

Wound care	Bandage removed at home by homecare staff (can shower but do not submerge) Negative pressure wound vac post-op 7 or 14 days post-op based on model
Assistive Device and Equipment	UH DME will provide pre-operatively or in the hospital Sling - Abduction sling should be worn for 4-6 weeks at surgeon's discretion
Home Health PT	If recommended, will be established with care transitions team prior to discharge (DC) home
Out-patient PT	Will complete following HHC for most patients. 3-4 weeks post-surgery date. Recommended to schedule prior to surgery. DO NOT DELAY SCHEDULING.

Total Shoulder Replacement Recommendations

Precautions (Strictly adhered to first 6 weeks post-op, guarded progression thereafter)

Depending on surgeon preference, limitations of flexion, scaption and external rotation may be recommended for at least the first 6 weeks post-operatively. Subscapularis repair is the main over-arching concern and range of motion (ROM) recommendations should be closely followed. Some surgeons may recommend a life-long restriction in active range of motion (AROM), as well as lifting, but specific restrictions should be discussed with surgeon at final follow-up visit.

Total Shoulder Replacement Yellow and Red Flags



YELLOW FLAGS

- Unhelpful beliefs about pain
- Expectations of poor outcomes
- Catastrophizing or fear of movement
- Over reliance on passive treatments



RED FLAGS -

Requires immediate referral for re-evaluation

- Severe or unrelenting pain
- Sudden inability to move the arm
- Unresolving numbness and tingling
- Excessive swelling
- Infection:
 - Fever, chills, night sweats
 - Redness, warmth around wound or incision
 - Abnormal pain at the surgical site
 - Drainage

ROM/Manual Therapy

- Early ROM as tolerated within the restricted ranges
- Soft tissue mobilization as needed, scar mobilization once incision heals (2-3 weeks)

Effusion

- Manage swelling and edema with cryotherapy and compression

Therapeutic Exercise

- Proper activation and recruitment of all shoulder and scapular musculature without compensation is required for successful strengthening
- Neuromuscular re-education is required for stabilization and correction of faulty mechanics

Functional Patient Outcomes

- Quick Disabilities of the Arm Shoulder and Hand (QuickDASH)
- Patient Specific Functional Scale (PSFS)

Considerations Regarding Running and Plyometrics

- Patients are advised to participate in low-impact exercise and activity
- Following surgery, goal is to prevent damage to new prosthetic
- High-impact activities are generally not advised

Patient should consult with physician if considering returning to running or other high impact sport.

Criteria for Discharge (Athletic)

- Physician clearance at last check-up
- Strength >90% compared to uninvolved lower extremity
- Demonstrate ability to simulate recreational activities
- Functional Patient Outcome Measures achieves MCID (minimum clinically important difference)

Criteria for Discharge (Non-Athletic)

- Criteria for discharge from PT is less rigorous for those not returning to sport
- Ensure the patient can perform all activities of daily living (ADLs) without pain, reactive effusion and with appropriate functional mechanics

Phase I Functional Mobility and Protection

TSA PHASE I: Day 0-2 Weeks

Precautions

- Abduction/traditional sling worn when not performing therapy exercises
- Monitor post-op incision healing for excessive bleeding/drainage or separation
- No AROM
- Non-weight bearing through affected upper extremity
- No IR or shoulder extension past neutral
- No reaching behind back
- *PROM limited to 90 degrees flexion and 30 degrees ER (scapular plane)
- Sleep posture: avoid operative side, discuss with your therapist

Goals

- Protection of implant
- Protect healing tissue and subscapularis
- Pain and swelling control
- Maintenance of cervical, elbow, and wrist/hand mobility/function
- Patient education regarding modification of ADLs to abide by precautions
- Gradual introduction of glenohumeral and scapulothoracic PROM

Interventions *(Exercise list is not exhaustive)*

Therapeutic Exercise / Activities / NMRE

- AROM of cervical spine, elbow, wrist
- PROM in scapular within above precautions
- Pendulums
- Table slides in flexion, forward bow
- Instruction in patient-performed PROM

Education: Activity Modification Modalities

- Cryotherapy
 - 5x daily for the 1st week
 - 3x daily for weeks 1-6

Criteria to Progress to Phase II

1. PROM scaption to 90 degrees
2. Independent with precautions and home exercise program (HEP)

*Criteria supersedes time for progression to next phase

*Dr. Victoroff: Limit A/PROM to 0 degrees 4+ weeks post-op. Consult with Dr. Victoroff as needed

Phase II Range of Motion, Isometrics, DC of Sling

PHASE II: 2 - 6 weeks

Precautions

- Continue using sling up to 4-6 weeks per surgeon discretion – off for exercise
- Avoid active IR and extension (reaching behind back)
- *Limit ER to 45 degrees with arm at side to protect subscapularis
- Avoid ER at 90 degrees abduction
- Passive elevation to 120 degrees in plane of the scapula
- Avoid supporting body weight through post-op UE
- Avoid lifting weighted objects heavier than 1-2 pounds, “no heavier than a coffee cup”

Goals

- Protect healing tissue
- Control pain and edema
- Prevent stiffness
- Protect subscapularis
- Continue to progress shoulder PROM
- Initiate active assisted range of motion (AAROM) and pain-free submax isometrics
- Regain independence with ADLs
- Consistent performance of HEP

Interventions *(Exercise list is not exhaustive)*

Therapeutic Exercise

- Continue elbow and wrist AROM
- Pendulums
- Scapular retraction
- Submaximal shoulder isometrics in neutral
 - Caution/avoid IR with subscap repair
- Gentle glenohumeral and scapulothoracic rhythmic stab

Range of Motion

- PROM/AAROM as tolerated within precautions
- AAROM c dowel – begin supine/reclined and progress to seated or standing
- Pulleys in scapular plane

Manual

- Soft tissue mobilizations
- Scar mobilization

Modalities

- **Cryotherapy:** continue as needed
Neuromuscular electrical stimulation (NMES) and/or Biofeedback as needed for recruitment

Criteria to Progress to Phase III

*Criteria supersedes time for progression to next phase

1. Good neuromuscular activation without compensation
2. Minimal pain or swelling
3. Shoulder scaption PROM to 120 degrees

Phase III Return to ADLs

PHASE III: 6- 12 weeks

Precautions

- No weight bearing through involved extremity
- No resisted IR x 12 weeks, no extension past neutral
- Passive ER not to exceed 60°
- No aggressive stretching - specifically IR behind the back
- No lifting greater than 5 pounds
- No quick or explosive movements

Goals

- Recruitment of scapulothoracic musculature and proper movement patterns
- Progress with isometrics and initiation of isotonic exercises
- Gradually improve shoulder AROM within precautions
- Out of sling
- Protect healing tissue, specifically subscapularis repair
- Progress from AAROM to AROM exercises
- Use of involved extremity for light ADLs - discomfort and stiffness may remain, or some pain with or after over-activity, etc.

Interventions (*Exercise list is not exhaustive*)

Range of Motion

- Continue as per Phase II as needed, Progress with PROM, light GH mobilization, no aggressive stretching

Therapeutic Exercises

- Initiate gentle IR stretching in scapular plane at 8 weeks and IR behind the back at 10 weeks
- Continue prior exercises, including isometrics and AAROM exercises
- Initiate AROM exercise progression from reclined to standing
- Scapular strengthening: rows, shoulder extension to neutral, prone exercises if appropriate, side lying ER
- Resisted deltoid exercises

Manual

- As needed

Modalities

- Cryotherapy, NMES, biofeedback and transcutaneous electrical nerve stimulation (TENS) as needed

Criteria to Progress to Phase IV

*Criteria supersedes time for progression to next phase

1. Achieve ~120° active flexion, ~ 60° ER, IR to sacrum
2. At least good (4+/5) UE strength or <15% deficit bilaterally

Phase IV Pain free ADLs and Return to Lifting

PHASE IV: 12 Weeks and Beyond

Precautions and Considerations for Return to Recreational Activities

This phase is only required for patients who wish to participate in moderate-impact recreational sports and/or return to work that involves heavy lifting, frequent lifting or frequent UE, shoulder and overhead movements.

- Patients who wish to return to moderate activities such as doubles tennis, doubles pickle ball or downhill skiing should only be on a case-by-case situation and should be cleared by their surgeon
- Jumping and contact sports are generally not recommended after a TSA; therefore, low to moderate impact activities are suggested in place of high impact sports
- No stretching limitations

Goals

- Correct compensatory movement patterns
- Ongoing interventions to improve neuromuscular control and proprioception
- Progressively return to recreational activities by increasing aerobic participation and restoring cardiovascular fitness
- Increase ROM and strength to align with patient activity level and goals - generally 150° flexion
- Progress towards DC to independent HEP addressing individual restrictions
- Initiate IR strengthening and closed chain exercises as appropriate
- Full return to all ADLs outside heavy overhead lifting

Interventions (*Exercise list is not exhaustive*)

Low Impact Activities

- Reactive motor control drills
- Progress core stability
- Stretching as needed to achieve goals
- Progression of previous strengthening exercises, including scapular, deltoid, rotator cuff and postural musculature
- Add progressive closed-chain strengthening
- Initiate IR strengthening exercises
- Advance cardio training (Bike, elliptical, row)

Therapeutic Exercise and Impact Progressions

- Progress from single to multiplanar movements with emphasis on neuromuscular control
- Sport-specific drills that fit low- to moderate-impact guidelines

Criteria to progress to moderate impact activities*

*Physician clearance required

1. Restore to functional, pain-free ROM
2. >85% strength compared to uninvolved UE
3. Patient demonstrates proper UE mechanics with functional testing and moderate impact activities

Total Shoulder Arthroplasty Rehabilitation Post-Operative Guidelines

Authors: Christopher LeVan, PT, John Campolieti, PT, Jeffrey Daniel Chelette, PT, Alison Ives, PT

Reviewed January 2025: Mario Gastaldo, PT

Physician Review February 2025: Daniel Zanotti, MD, Robert Zanotti, MD, Brian Victoroff, MD, Jacob Calcei, MD, John Martin Leland, MD

References:

Baumgarten KM, Osborn R, Schweinle WE 3rd, et al. The position of sling immobilization influences the outcomes of anatomic total shoulder arthroplasty: a randomized, single-blind, prospective study. *J Shoulder Elbow Surg* 2018;27(12): 2120–8.

Bullock GS, Garrigues GE, Ledbetter L, Kennedy J. A Systematic Review of Proposed Rehabilitation Guidelines Following Anatomic and Reverse Shoulder Arthroplasty. *Journal of Orthopaedic and Sports Physical Therapy*. 2019; 49(5): 337-346.

Ellenbecker TS, Manske RC, Kelley M. The shoulder: physical therapy patient management using current evidence. In: Hughes C, ed. *Current Concepts of Orthopedic Physical Therapy*. 4th Ed. La Crosse, WI. Academy of Orthopedic Physical Therapy; 2016.

Gillespie RJ, Garrigues GE, Chang ES, et al. Surgical exposure for reverse total shoulder arthroplasty: differences in approaches and outcomes. *Orthop Clin North Am*. 2015 Jan;46(1):49-56. doi: 10.1016/j.ocl.2014.09.015. PMID: 25435034.

Howard MC, Trasolini NA, Waterman BR. Optimizing outcomes after reverse total shoulder arthroplasty: rehabilitation, expected outcomes, and maximizing return to activities. *Current Reviews in Musculoskeletal Medicine*. 2023; 145-153.

Kennedy JS, Garrigues GE, Pozzi F, et al. The American Society of Shoulder and Elbow Therapists' consensus statement on rehabilitation for anatomic total shoulder arthroplasty. *J Shoulder Elbow Surg*. 2020; 29(10): 2149-2162.

Levy JC, Everding NG, Gil CC, et al. Speed of recovery after shoulder arthroplasty: a comparison of reverse and anatomic total shoulder arthroplasty. *J Shoulder Elbow Surg*. 2014;23(12):1872-1881. doi: 10.1016/j.jse.2014.04.014. Epub 2014 Jun 26.

Molé D, Wein F, Dézaly C, et al. Surgical technique: the anterosuperior approach for reverse shoulder arthroplasty. *Clin Orthop Relat Res*. 2011;469(9):2461-2468. doi: 10.1007/s11999-011-1861-7.

Pathak A, Wilson R, Sharma S et al. Measurement properties of the patient-specific functional scale and its current uses: an updated systematic review of 57 studies using COSMIN guidelines. *J Orthop Sports Phys Ther*. 2022;52(5):262–275. doi: 10.2519/jospt.2022.10727. Epub 2022 Feb 05.

Seitz AL, Christain H, Lutz A, Shanley E. The shoulder: evidence-informed physical therapy patient management. In: Simoneau G, ed. *Current Concepts of Orthopedic Physical Therapy*. 5th Ed. La Crosse, WI. Academy of Orthopedic Physical Therapy; 2021.

Schick S et al. Variability in physical therapy protocols following total shoulder arthroplasty. *Clin Shoulder Elbow*. 2023; 26(3): 267-275.

The flag system. Physiopedia. Updated 2024. Accessed October 4, 2024. https://www.physio-pedia.com/The_Flag_System.